

## Unravelling the evolutionary history and future prospects of endemic species restricted to former glacial refugia

ORLY RAZGOUR,<sup>\*1</sup> IRENE SALICINI,<sup>†</sup> CARLOS IBÁÑEZ,<sup>†</sup> ETTORE RANDI<sup>‡ §</sup> and JAVIER JUSTE<sup>†</sup>

<sup>\*</sup>Division of Biological and Environmental Sciences, School of Natural Sciences, University of Stirling, Stirling, Scotland FK9 4LA, UK, <sup>†</sup>Estación Biológica de Doñana (CSIC), Avda Americo Vespucio s/n, 41092 Seville, Spain, <sup>‡</sup>Laboratorio di Genetica, Istituto Superiore per la Protezione e Ricerca Ambientale, Via Ca' Fornacetta 9, Ozzano dell' Emilia, Bologna, Italy,

<sup>§</sup>Department 18/ Section of Environmental Engineering, Aalborg University, Sohngårdsholmsvej 57, 9000 Aalborg, Denmark

### Abstract

The contemporary distribution and genetic composition of biodiversity bear a signature of species' evolutionary histories and the effects of past climatic oscillations. For many European species, the Mediterranean peninsulas of Iberia, Italy and the Balkans acted as glacial refugia and the source of range recolonization, and as a result, they contain disproportionately high levels of diversity. As these areas are particularly threatened by future climate change, it is important to understand how past climatic changes affected their biodiversity. We use an integrated approach, combining markers with different evolutionary rates and combining phylogenetic analysis with approximate Bayesian computation and species distribution modelling across temporal scales. We relate phylogeographic processes to patterns of genetic variation in *Myotis escalerai*, a bat species endemic to the Iberian Peninsula. We found a distinct population structure at the mitochondrial level with a strong geographic signature, indicating lineage divergence into separate glacial refugia within the Iberian refugium. However, microsatellite markers suggest higher levels of gene flow resulting in more limited structure at recent time frames. The evolutionary history of *M. escalerai* was shaped by the effects of climatic oscillations and changes in forest cover and composition, while its future is threatened by climatically induced range contractions and the role of ecological barriers due to competition interactions in restricting its distribution. This study warns that Mediterranean peninsulas, which provided refuge for European biodiversity during past glaciation events, may become a trap for limited dispersal and ecologically limited endemic species under future climate change, resulting in loss of entire lineages.

**Keywords:** approximate Bayesian computation, bats, climate change, *Myotis escalerai*, phylogeography, species distribution modelling

Received 6 October 2014; revision received 3 September 2015; accepted 4 September 2015

### Introduction

The contemporary distribution and genetic composition of biodiversity bear a signature of species' evolutionary

Correspondence: Orly Razgour, Fax: +44 1786 467843; E-mail: Orly.Razgour@gmail.com

<sup>1</sup>Present address: School of Biological Sciences, University of Bristol, Life Sciences Building, Bristol BS8 1TQ, UK

histories. Quaternary climatic oscillations, in the form of recurring glacial–interglacial cycles, resulted in substantial range shifts, population extinctions and lineage divergences (Hewitt 2000), although effects varied with latitude, topography (Hewitt 2004) and individual species' adaptations and environmental tolerances (Stewart *et al.* 2010).

With the advent of molecular tools, the study of the distribution of biodiversity was extended to include

genetic relationships between individuals and the influence of historical processes on the geographic distribution of genetic lineages (Avise 2000). Phylogeography has provided the framework to determine the causal links between geography, climate change, ecological interactions and the evolution of taxa (Hickerson *et al.* 2010). Its integration with ecological niche modelling has helped elucidate the processes and mechanisms shaping genetic variation and the evolutionary trajectories of species and populations (Alvarado-Serrano & Knowles 2014). Understanding the phylogeographic structure of species, and the mechanisms that sustain it, is integral to conserving their full genetic diversity and to managing evolutionary significant units within species according to their differing regional vulnerabilities (Schmitt 2007). Moreover, understanding species' responses to past events may help us better predict the potential consequences of future climatic changes (Hofreiter & Stewart 2009).

During Pleistocene glacial periods, much of northern and central Europe was covered by ice sheets and permafrost. The Mediterranean peninsulas of Iberia, Italy and the Balkans acted as glacial refugia for many European species and as the source of rapid northern range colonization during interglacial, warmer climatic periods. Cycles of contraction-expansion into and out of glacial refugia resulted in a genetic signature of southern richness with deep divergence between refugial populations vs. northern impoverishment and genetic homogeneity (Hewitt 2004). Stable areas that persisted across glaciation cycles harbour particularly high levels of species richness (Araújo *et al.* 2008) and unique genetic diversity (Hampe & Petit 2005), and as a result are of high evolutionary importance (Stewart *et al.* 2010). However, these hotspots of genetic diversity are particularly threatened by future climate change (EEA 2012; Razgour *et al.* 2013), and therefore, it is important to understand how past climatic changes affected their biodiversity.

The Iberian Peninsula has a rich and well-studied biogeographic history. Its complex topography and geographic position between the Mediterranean and North Atlantic creates distinct bioclimatic regions with ecologically and genetically divergent taxa (Gómez & Lunt 2007). Yet, this great environmental heterogeneity, combined with relative climatic stability and long-term lineage persistence and divergence without large geographic displacement, makes it more difficult to interpret the genetic population structure and evolutionary history of species within Iberia (Hewitt 2001; Rodriguez-Sánchez *et al.* 2010). The Iberian Peninsula played an important role in the evolutionary history of European bats. Phylogeographic studies of widely distributed European bat species show that although Iberia


was an important glacial refugium for many species, in some cases, it did not necessarily contribute to post-glacial range recolonization because lineages remained isolated inside the peninsula by the Pyrenees mountain range (e.g. *Barbastella barbastellus*, Rebelo *et al.* 2012; and *Rhinolophus hipposideros*, Dool *et al.* 2013). However, for other bat species, the Iberian refugium was the main source of range recolonization, while the Alps formed a stronger barrier to range expansion from other Mediterranean refugia (e.g. *Myotis myotis*, Ruedi & Castella 2003).

Here, we set to unravel the effect of Quaternary climatic oscillations on the evolutionary history of *Myotis escalerae*, a bat species endemic to the Iberian Peninsula (defined here as the area including Spain, Portugal, the Balearic Islands, Andorra and the French Pyrenees), and to determine factors that limit its distribution and how it will be affected by future climate change. *M. escalerae* is part of the *Myotis nattereri* cryptic species complex (*M. nattereri sensu stricto*, *M. escalerae*, *M. spA* and *M. spB*; Salicini *et al.* 2011) that has only recently been genetically confirmed as a separated species (Ibáñez *et al.* 2006), but has been described morphologically more than a century ago (Cabrera 1904). Unlike other bat species, the entire evolutionary history of *M. escalerae* took place within Iberia (Salicini *et al.* 2013), and therefore, both its present genetic population structure and future survival are closely linked to climate change processes within the Iberian Peninsula. We use an integrated approach, combining markers with different evolutionary rates, and combining phylogenetic analysis with approximate Bayesian computation (ABC) model-based inference and species distribution modelling (SDM) across temporal scales, to relate phylogeographic processes to contemporary and future patterns of genetic variation.

## Methods

### Sample collection

Genetic samples, in the form of 3-mm wing biopsies, were collected from 252 *M. escalerae* bats captured in 16 colonies, located mostly in underground sites (caves), distributed across the Iberian Peninsula and the Balearic Island of Mallorca (Table S1, Supporting information, Fig. 1A). Sequences of two individuals from the Sevilla colony were previously used in Salicini *et al.* (2011, 2013) (GenBank Accession nos: JN591489.1 and JX826314.1). In addition to the samples from the 16 colonies, the mitochondrial DNA analysis also included *M. escalerae* sequences downloaded from GenBank that belonged to samples from the French Pyrenees (JF412390 and JF412391—Puechmaille *et al.* 2012) and



**Fig. 1** *Myotis escalerai* population structure based on the mtDNA (cytochrome *b*) data set (colonies and individual samples,  $N = 359$ ; Tables S1–S2, Supporting information). (A) Map of the location of the colonies (larger circles and names) and individual samples included in the study, colour coded based on phylogenetic clades. (B) Bayesian phylogenetic tree showing posterior probability values  $>0.8$ . Haplotypes are named based on their respective sampling locations, and clades are marked with their respective colours (yellow—west, blue—south and red—north-central-east). (C) Median-joining haplotype network, colour coded based on phylogenetic clades. Circle sizes correspond to number of samples. Numbers indicate haplotypes separated by  $>1$  mutation. Haplotype CasGiB is marked with A, and LROurSeg with B.

108 *M. escalerai* sequences from across the Iberian Peninsula, including 21 samples from around the Pyrenees (Navarra, Huesca, Lleida and Girona) and 18 samples from adjacent areas (La Rioja, Zaragoza, Teruel and Tarragona). Out of the individual sequences, 24 were taken from previous studies (Salicini *et al.* 2011, 2013). These samples were added to better characterize the range of the species and to provide better coverage of areas of sympatry with *M. spA* (Table S2, Supporting information; Fig. 1A).

#### Laboratory procedures

Genomic DNA was extracted from all samples following the methods described in Salicini *et al.* (2013). We sequenced 750 bp of the mitochondrial DNA (mtDNA) cytochrome *b* (*Cyt b*) gene, using the primers Molcit-F (Ibáñez *et al.* 2006) and Molcit-R (Salicini *et al.* 2011). PCR conditions and sequencing information are outlined in Salicini *et al.* (2013). Sequences were aligned and edited using SEQUENCHER 4.5 (Gene Codes Corp, MI, USA) and collapsed into unique haplotypes with DAMBE v5.2.31 (Xia & Xie 2001).

Samples were genotyped for 10 microsatellite loci previously published for the genus (A13, D9, D15, E24, F19, G25, H29, A24, H23; Castella & Ruedi 2000; and b22; Kerth *et al.* 2002). The forward primer of each locus pair was labelled fluorescently with HEX or 6-FAM (Applied Biosystems). Microsatellites were combined into single or double PCR sets. Each PCR mix contained 0.3 µl primer sets at 10 µM, 1 µl of PCR buffer 10×, 0.3 µl dNTPs, 0.05 µl TAQ and 1 µl of DNA, adding H<sub>2</sub>O up to 10 µl total volume. When needed, 0.8 µl of bovine serum albumin was added. PCR amplification was performed using ABI Veriti thermal cycler (Applied Biosystems, USA). We used the following PCR program: initial denaturation at 95 °C for 5 min, followed by 30–40 cycles of 95 °C for 30 s, annealing temperature from 55 °C to 60 °C, depending on the primers, for 45 s and 72 °C for 45 s, followed by a final extension at 72 °C for 10 min. PCR products were sequenced using ABI 3130 48-well DNA Sequencer. Allele sizes were assigned using the GENEMAPPER software (Applied Biosystems, USA).

Observed and expected heterozygosity and estimated null allele frequencies were calculated using CERVUS v3.0.3 (Kalinowski *et al.* 2007) and MICRO-CHECKER (Van Oosterhout *et al.* 2004). Tests for departures from Hardy–Weinberg equilibrium and assessment of linkage disequilibrium were performed in GENEPOL v4.0.10 (Raymond & Rousset 1995; Rousset 2008). Loci that were out of Hardy–Weinberg equilibrium and with a high frequency of null alleles in several populations were removed from the analysis.

### Genetic data analysis

*Mitochondrial data set.* We used JMODELTEST v2.1.6 (Darriba *et al.* 2012) to select the Hasegawa–Kishino–Yano (HKY) mtDNA substitution model with gamma-distributed rate variation based on the Bayesian information criterion (BIC) values. Bayesian phylogenetic trees were constructed in MRBAYES v3.2.1 (Ronquist & Huelsenbeck 2003), using two *Myotis spA* sequences as outgroup to root the tree. We ran  $5 \times 10^7$  generations with four chains, sampled every 500th generation, and two simultaneous runs, discarding the first 25% of trees generated as burn-in. Trees and posterior probabilities were visualized with FIGTREE v1.3.1 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Parsimony haplotype network was constructed with NETWORK (v4.610; Fluxus Technology), employing the median-joining network algorithm and the Greedy FHP distance calculation method. Nucleotide polymorphism, haplotype diversity, genetic divergence and differentiation between populations were calculated in DNASP v5.10 (Librado & Rozas 2009), with 10 000 permutations to obtain probability values.

*Microsatellite data set.* Analysis of microsatellite genetic diversity, including allele frequencies, number of private alleles, allelic richness, heterozygosity, gene diversity and population differentiation ( $F_{ST}$ ), was carried out at the colony level with GENALEX v6.4 (Peakall & Smouse 2006) and FSTAT v2.9.3.2 (Goudet 1995) controlling for differences in sample sizes. To test for levels of relatedness among individuals, we used the triadic maximum likelihood estimator (TrioML; Wang 2007) implemented in Coancestry (Wang 2011) because this measure allows for inbreeding and accounts for genotyping errors in the data.

Population structure in the microsatellite data set was inferred using individual-based Bayesian assignment tests implemented in STRUCTURE v2.3.3 (Pritchard *et al.* 2000). Number of tested genetic clusters (K) ranged from 1 to 15. We performed 10 independent runs for each K, using the general admixture model with correlated allele frequencies and  $10^6$  Markov chain Monte Carlo (MCMC) generations following a burn-in phase of  $5 \times 10^5$  generations. The number of distinct clusters was determined using STRUCTURE HARVESTER (Earl & von Holdt 2012) based on the number of clusters at which the mean log-likelihood peaked and where variation among runs was minimal. Cluster assignment was visualized with DISTRUCT (Rosenberg 2004).

Because the presence of closely related individuals (in particular full-siblings) can bias the number of clusters identified in the STRUCTURE analysis (Rodriguez-Ramilo

& Wang 2012), we first ran assignment tests with the whole data set and then reran the analysis removing individuals with TrioML values  $>0.5$ . This threshold was selected because below this value, most of the pairwise estimations were among individuals from geographically distant colonies (58% for TrioML = 0.5 vs. 2% for TrioML  $>0.5$ ).

### SDM procedures

We used SDMs to generate phylogeographic hypotheses for testing with ABC inference, to identify environmentally stable areas where the species persisted overtime, to determine the most important environmental variables limiting the distribution of *M. escalerai* and to predict future changes to distribution (Alvarado-Serrano & Knowles 2014). We predicted the potential distribution of suitable conditions for *M. escalerai* under present, past (LGM ~21 000 years before present and the last interglacial period [LIG] ~130 000 ybp) and future (2070) climatic conditions. Study area extent was set as the Iberian Peninsula, the Balearic Islands and France up to latitude 49.5 N and longitude 6.5 E. This extent enabled the inclusion of potentially suitable areas beyond the species' currently known distribution, while limiting problems associated with selecting pseudo-absences at large distances from known location records (VanDerWal *et al.* 2009). Model resolution was ~1 km (30 arc seconds).

Models were generated with MAXENT v3.3.3 (Phillips *et al.* 2006) using 135 location records, including one location record from the French Pyrenees taken from Evin *et al.* (2009). As the whole of Iberia has been sampled extensively for this species, our data set is not likely to suffer from sampling bias. All location records were genetically confirmed because of potential range overlap with cryptic congeners of the *M. nattereri* species complex. We used the Average Nearest Neighbor tool in ARCGIS v10.2 (ESRI) to remove duplicate and clustered location records to minimize spatial autocorrelation between location records.

We ran two types of models, climatic-topographic models (climate model) for all time periods, and a full model that included also habitat variables for the present only. Climatic and topographic layers were downloaded from WORLDCLIM (<http://www.worldclim.org>), geological layers from One Geology (<http://www.onegeology.org/>, reclassified into 18 broad categories) or USF Geoportal Data Depository (Karst Regions of the World, <http://gisdata.rc.usf.edu/>, Hollingsworth *et al.* 2008). Habitat variables were obtained from the European Space Agency (GlobCover 2009, [http://due.esrin.esa.int/page\\_globcover.php](http://due.esrin.esa.int/page_globcover.php)) for land cover (reclassified into 10 categories), European

Environment Agency (Corine Land Cover 2006, <http://www.eea.europa.eu/>) for woodland variables (woodland type and distance to woodlands) and Hansen *et al.* (2013) for per cent tree canopy cover. Multicollinearity among environmental variables was tested with ENMTOOLS v1.3 (Warren *et al.* 2010), removing highly correlated variables (correlation coefficients  $\geq 0.8$ ) and variables that did not contribute to the SDMs. The following layers were included in the final models: maximum temperature of warmest month (BIO5), minimum temperature of the coldest month (BIO6), average temperature of the driest quarter (BIO9), temperature seasonality (BIO4), annual rainfall (BIO12), rainfall seasonality (BIO15), rainfall in warmest quarter (BIO18), slope, altitude, distance to karsts (maternity colonies are known to form in caves; Ibáñez *et al.* 2006), land cover, distance to woodlands and per cent tree cover.

Models were projected into the past using the CCSM and MIRCO general circulation models (GCMs) for the LGM and one LIG model. Future models for 2070 were generated using three European GCMs (HadGEM2\_ES, IPSL-CM5A-LR and MPI-ESM-LR), and the IPCC5 +8.5 W/m<sup>2</sup> Representative Concentration Pathways (IPCC 2013), representing the 'worst case' scenario.

Our modelling procedures followed recommendations in Merow *et al.* (2013). We compared several models with different variables and parameter combinations (regularization values, number of features included) in ENMTOOLS and selected the best models based on Akaike information criterion (AIC) scores. The final full and climate models included all features with a regularization value of 1, 10 000 background points and 1500 iterations. When comparing models, we used the raw output, but when running the final models, we used the cumulative output. Projected output maps generated by the different LGM or future GCMs were multiplied to produce a single map per time period. To determine differences in the extent of areas with high relative occurrence probability over time, we converted model outputs into binary maps using the thresholding method that maximizes the sum of sensitivity and specificity (Liu *et al.* 2013). Range changes were calculated for the Iberian Peninsula alone (including the Pyrenees). Maps were processed in ARCGIS v10.2 (ESRI).

Model predictive ability was tested with fivefold cross-validation and compared based on the area under the curve (AUC) of the receiver operator characteristics. To determine whether our models performed significantly better than random, we tested whether our models' training and test AUC scores fell outside the 95% confidence intervals of the distribution of the AUC scores of 100 null models (Raes & ter Steege 2007), randomly generated in ENMTOOLS with the altitude layer.

### ABC framework

The evolutionary history of *M. escalerai* was reconstructed using the ABC approach implemented in DIYABC v2.0.4 (Cornuet *et al.* 2014) to identify source populations and patterns of colonization. Phylogeographic hypotheses were generated based on paleo-SDM predictions. We first ran a full analysis (Analysis 1), which included all colonies, divided into three geographical groups (western, southern and north-central-eastern). The full analysis aimed to identify the source population, LGM refugial populations and patterns of post-LGM range recolonization. Next, we ran separate ABC analyses for the geographically separated western (Analysis 2) and eastern (Analysis 3) groups to identify the representative putative source colonies of each group in relation to predicted climatic suitability during the LGM. Finally, in Analysis 4, we assessed the demographic history of the western and eastern groups, comparing scenarios of post-LGM population expansion vs. pre-/post-LGM population declines (Fig. S8, Supporting information). Scenarios compared in each analysis and their specific demographic parameters are outlined in Supplementary Materials.

Approximate Bayesian computation analyses were carried out with the combined microsatellite and mtDNA data sets as well as on each data set separately. The separate mtDNA analysis also included the 108 individual samples and the two French sequences from GenBank. The remaining analyses only included samples from the 16 colonies. We generated  $10^6$  simulations for each scenario tested in each analysis. The posterior probability of scenarios was estimated using a weighted polychotomous logistic regression. We checked model performance and empirically evaluated the power of the model to discriminate among scenarios (confidence in scenario choice) by simulating pseudo-observed data sets with the different scenarios and calculating false allocation rates (type 1 and 2 errors, Cornuet *et al.* 2010).

## Results

### Mitochondrial DNA data set

We identified 50 unique Cyt b haplotypes (20 from the 16 colonies). The Bayesian phylogenetic tree showed maximum posterior probability support for the split of *M. escalerai* into two principal lineages, the western (southwest clade in Salicini *et al.* 2013) and southern clade, and the remaining haplotypes, which mainly constituted of north-central-eastern haplotypes. The western and southern clades were further divided (posterior probability = 0.85) into the western and southern lineages (Fig. 1B).

Similarly, the haplotype network divided the haplotypes into three separate haplogroups: western, southern and north-central-eastern. Western haplotypes were separated from the remaining haplotypes by >19 mutational steps (per cent differences >2.5%). Most southern haplotypes grouped together and were separated by >8 mutational steps from the north-central-eastern haplotypes (>1.1%). However, one haplotype from the southeastern colony Granada grouped with the north-central-eastern haplotypes, while most of the samples from the southern colony of Sevilla grouped with the western haplogroup. Samples from the French Pyrenees belonged to the common eastern haplotype (CasGiB), as did most samples from around the Pyrenees. However, some unique haplotypes were identified in the Pyrenees, all of which were separated by one mutational step from either the common eastern (CasGiB) or north-central (LROurSeg) haplotypes, depending on their geographical location (eastern and central vs. western Pyrenees) (Fig. 1C, Table S1–S2, Supporting information).

Mitochondrial haplotype diversity was highest in the north-central-eastern group, even after accounting for differences in sample size (32 haplotypes, 0.16 per sample), but nucleotide diversity was highest in the southern group ( $\pi = 0.016$ ; Table S3, Supporting information). Among the colonies, Cádiz and Illes Balears had the highest haplotype diversity, while Granada, Alacant and Sevilla the highest nucleotide diversity (Table 1). Overall genetic differentiation at the mtDNA level between the western, north-central-eastern and southern geographic groups was significant ( $\chi^2_{90} = 678.5$ ,  $P < 0.001$ ; overall  $\theta_{ST} = 0.73$ ), with particularly high levels of differentiation between the western and north-central-eastern groups ( $\theta_{ST} = 0.93$ ; Table S4, Supporting information).

#### *Microsatellite data*

Of the 10 microsatellite loci, one marker (H29) was removed due to high frequency of null alleles. After removing this marker, all colonies, but Huelva, were overall in Hardy–Weinberg equilibrium. None of the markers were in linkage disequilibrium and all were in Hardy–Weinberg equilibrium in at least 13 of the 16 colonies. The data set, excluding H29, contained a total of 103 alleles, with an average of  $11.44 \pm 5.5$  alleles per locus (range 4–21), and 10 private alleles.

Genetic diversity (adjusted for sample size) in terms of allelic richness, heterozygosity, gene diversity and number of private alleles was highest in Granada (southern Iberia) followed by Cáceres (western Iberia) and was lowest in Girona (eastern Iberia) and Illes Balears (Table 1). Levels of relatedness were particu-

larly high within the Girona and Illes Balears colonies (mean TrioML =  $0.44 \pm 0.1$  and  $0.25 \pm 0.2$ , respectively), whereby a third of the pairwise relatedness values between individuals within the Girona colony were >0.5. Levels of population differentiation were highest between Girona and all other colonies and Illes Balears and all other colonies, even after the removal of close relatives. Particularly low levels of differentiation were found between Cáceres and most southern and western colonies and among the southern colonies (Table S5, Supporting information).


Individual-based Bayesian assignment tests detected genetic population structure in *M. escalerai*. Individuals were best divided into four genetic clusters (Ln probability ( $K$ ) =  $-7730 \pm 5$ ; Fig. S1, Supporting information), despite some level of admixture in most colonies. The most northeastern colony, Girona, formed a separate cluster; however, this cluster disappeared once close relatives (TrioML >0.5) were removed from the analysis. Individuals whose haplotypes belonged to the mtDNA north-central-eastern clade tended to be assigned to different clusters from individuals from the mtDNA western clade, with the exception of individuals from the most northwestern colony (Ourense). However, most individuals whose haplotypes belonged to the mtDNA southern clade showed high levels of admixture between clusters, and only an east to west geographic gradient was evident at the nuclear microsatellite level (Fig. 2).

#### *SDM across temporal scales*

All SDMs had high predictive ability, did not overfit presence data (full model: AUC = 0.89 AUC<sub>cross-validation</sub> =  $0.80 \pm 0.04$ ; climatic model: AUC = 0.87, AUC<sub>cross-validation</sub> =  $0.79 \pm 0.03$ ) and had significantly higher predictive ability than the null models (mean AUC =  $0.64 \pm 0.004$  [95% confidence intervals], range: 0.57–0.67). The best fit model in terms of AIC scores had a regularization value of 1. The main ecogeographical variable contributing to both the climatic and full models was slope. Other important variables contributing to the climatic model were annual rainfall (BIO12), temperature seasonality (BIO4), rainfall seasonality (BIO15) and average temperature of the dry quarter (BIO9), while the habitat variable per cent tree cover and the land cover type conifer woodlands was important in the full model (Figs S2–S3, Supporting information). Both models show high concordance on predictions for areas occupied by the 16 colonies, although the full model offers a finer resolution, which results in more fragmented habitat suitability in the northwest. All colonies, except for two western colonies (Nabão and Ourense), are currently located in areas

**Table 1** Genetic diversity of *Myotis escalerai* colonies based on the microsatellite (first six columns) and the mitochondrial DNA (last four columns) data sets, with sample sizes presented in brackets. Mean allelic richness and gene diversity ( $\pm$ standard deviation) were adjusted based on sample size

	Mean number of alleles	Shannon Index	Gene diversity	Allelic richness	Number of private alleles	Heterozygosity ( $H_e$ )	Number of haplotypes	Number of polymorphic sites	Haplotypic diversity	Nucleotide diversity ( $P_i$ )
Alacant (19)	7.44 ± 1.0	1.593	0.76 ± 0.1	6.0 ± 2.2	0	0.739	3	8	0.608	0.0044
Amarela (11)	5.00 ± 0.5	1.261	0.67 ± 0.1	4.9 ± 1.5	0	0.636	1	0	0	0
Cáceres (17)	7.33 ± 1.1	1.609	0.75 ± 0.2	6.3 ± 2.6	3	0.728	1	0	0	0
Cádiz (11)	6.33 ± 0.7	1.579	0.77 ± 0.2	6.2 ± 2.1	0	0.735	4	3	0.691	0.0014
Castellón (19)	7.22 ± 1.0	1.586	0.74 ± 0.2	6.1 ± 2.3	0	0.719	1	0	0	0
Entrimio (16)	6.33 ± 0.7	1.433	0.69 ± 0.2	5.6 ± 1.8	0	0.672	1	0	0	0
Girona (17)	3.44 ± 0.7	0.792	0.44 ± 0.3	3.1 ± 1.6	0	0.431	1	0	0	0
Granada (18)	8.11 ± 1.2	1.711	0.77 ± 0.2	6.8 ± 2.9	4	0.748	2	10	0.526	0.0070
Huelva (16)	6.00 ± 0.7	1.440	0.70 ± 0.2	5.4 ± 2.0	0	0.681	1	0	0	0
I. Balears (16)	5.44 ± 0.9	1.192	0.60 ± 0.2	4.7 ± 2.2	2	0.581	4	3	0.792	0.0019
La Rioja (18)	6.00 ± 0.9	1.396	0.70 ± 0.2	5.2 ± 2.0	1	0.681	2	1	0.118	0.0002
Nabão (11)	5.78 ± 0.7	1.474	0.74 ± 0.2	5.6 ± 2.0	0	0.702	2	3	0.327	0.0013
Ourense (15)	6.22 ± 0.9	1.433	0.69 ± 0.2	5.6 ± 2.4	0	0.666	1	0	0	0
Preguiça (14)	5.67 ± 0.6	1.305	0.65 ± 0.2	5.2 ± 1.5	0	0.622	3	4	0.538	0.0012
Segovia (17)	7.44 ± 1.1	1.568	0.72 ± 0.2	6.3 ± 2.6	0	0.700	1	0	0	0
Sevilla (17)	7.56 ± 1.0	1.600	0.75 ± 0.1	6.3 ± 2.2	0	0.728	2	25	0.111	0.0037




**Fig. 2** *Myotis escalerai* population structure based on the microsatellite data set. (A) STRUCTURE analysis including all samples ( $K = 4$ ); and (B) STRUCTURE analysis after close relatives ( $\text{TrioML} > 0.5$ ) was removed ( $K = 3$ ), showing cluster membership plots and frequency of each cluster in the studied colonies.

predicted to have a high relative occurrence probability for *M. escalerai*, although both are still within 5 km distance of suitable areas (Fig. 3A–B; Fig. S4, Supporting information).

Paleo-SDMs predicted a substantial decrease in the extent of suitable conditions for *M. escalerai* in Iberia during the LGM compared with present conditions (per cent of area above suitability threshold for present: 34%, for LGM: 8.4%). Suitable climatic conditions during the LGM were restricted to isolated areas in the central west, south and east of Iberia and in southeastern France, while the central Plateaus, western Pyrenees and the north and west coasts were climatically unsuitable. As a result, in the western Group, only the most central colony Cáceres and the northern colony Entrimio were located in climatically suitable areas (Fig. 3C–D). Model predictions were affected by vari-

ables outside their training range around the Pyrenees, northwest Iberian coast and northern France (Fig. S6, Supporting information). The extent of suitable conditions was also low during the LIG (17%), but suitable areas were restricted to the north Atlantic coast, western Iberia (Portugal) and the southern tip near the Strait of Gibraltar (Fig. S5, Supporting information).

Future SDMs predicted a reduction in range suitability for *M. escalerai* in Iberia by 2070 (to 18.1%) with most of the south and interior of Iberia predicted to become climatically unsuitable. However, the northern Atlantic coast, Pyrenees and northwestern France are predicted to gain suitable areas (Fig. 3E–F). This will result in the majority of colonies and the entire southern lineage being located in climatically unsuitable areas by 2070. However, these predictions should be considered with caution because temperature variables



**Fig. 3** Species distribution models for *Myotis escalerai* across temporal scales: (A–B) present climate model, (C–D) last glacial maximum (LGM ~21 000 ybp) and (E–F) future (2070, +8.5rcp scenario). Models are presented as a scale of relative occurrence probability from low in yellow to high in dark blue (A, C, E) or as binary maps of potentially suitable areas in black (B, D, F). White circles denote the location of the studied colonies.

were outside their training range across most of the peninsula (Fig. S7, Supporting information).


#### ABC inference of demographic/evolutionary history

Model-based inference pointed to the western group being the source population of *M. escalerai* and to the presence of two separate refugia in the Iberian Peninsula during the LGM, one in the west and one in the north-central-east. The southern population diverged from the western population after the end of the LGM, and later was admixed with gene flow from the north-central-east population (Scenario 1.1, posterior probability = 0.93; type 1 error = 0.03, type 2 = 0.02; Fig. 4A; Table S6, Supporting information).

The western group analysis identified Cáceres as the representative source population of the western *M. escalerai* group, from which all other colonies split

after the LGM, beginning with the most southwestern colony (Amarela) and ending with the adjacent central colony (Nabão) (Scenario 2.1, posterior probability = 0.99; type 1 error = 0.02, type 2 error = 0.02; Fig. 4B; Table S7, Supporting information). Similarly, Castellón was the representative source population in the best supported model for the eastern group and all other eastern colonies split directly from this population post-LGM, with the oldest split being between Castellón and Girona (Scenario 3.1, posterior probability = 0.83; type 1 = 0.05, type 2 = 0.03; Fig. 4B; Table S8, Supporting information). In both analyses, population split dates were estimated to have occurred between the early and mid-Holocene.

Demographic history modelling indicates that the western group's effective population size has increased more than 10-fold after the end of the LGM, while the eastern population size remained stable, although cur-



**Fig. 4** Results of the approximate Bayesian computation analysis of the evolutionary history of *Myotis escalerai*, showing the selected scenarios for (A) the full model and (B) the western and eastern group analyses. White circles denote the location of colonies. Arrows represent the direction of colonization from the source population, with median estimated divergence dates.

rently both groups have similar estimated effective population sizes (Scenario 4.3, posterior probability = 1.0, type 1 = 0.016, type 2 = 0.01; Fig. S8, Supporting information; Table S9, Supporting information). The timing of the western population expansion corresponds with the estimated time of colonization of the southwestern colonies and therefore may reflect population expansion to areas south of the LGM refugia.

The same full model scenario (Scenario 1.1) was supported by the microsatellite-only data set (posterior probability = 0.99, type 1 = 0.05, type 2 = 0.06). There was no clear support for models ran using the extended mtDNA-only data set, which included the 16 colonies and all the individual samples. Although Scenario 1.4 (west source, colonized east via south) received relatively high support (posterior probability = 0.73), error rates were high (type 1 = 0.39, type 2 = 0.32), indicating that the analysis was unable to differentiate between the scenarios.

## Discussion

The combination of climate change and topographically originated environmental heterogeneity played an important role in shaping the evolutionary history and current genetic population structure of *M. escalerai* within the Iberian refugium, and it is likely to continue shaping the future distribution and patterns of genetic diversity of this restricted range endemic species.

### *The biogeographic history of Myotis escalerai*

It is not clear what event caused the divergence of *M. escalerai* from its Moroccan cryptic sister species *M. spB* around 0.99 million years ago (Salicini *et al.* 2013). However, despite this relatively recent speciation event, we found strong support for divergence into distinct clades. Quaternary climatic oscillations appear to have left a signature of geographic population structure in *M. escalerai* which corresponds to patterns of deep lineage divergence in other Iberian taxa whose lineages diverged before the Pleistocene (e.g. the *Vipera latastei/monticola* group, Velo-Anton *et al.* 2012).

Based on the mtDNA data set, *M. escalerai* across Iberia is divided into three main lineages, the western clade, which is restricted to the Atlantic climatic region in Portugal, the north-central-eastern clade and the southern clade. Paleo-SDMs indicate that this split may be the result of the disjunct distribution of suitable climatic conditions during the LGM when suitable areas were restricted to isolated patches in the west, east, south and near the Pyrenees and southern France. Model projections and the strong association of the phylogenetic divide with geography lend support to the

suggestion that during the Pleistocene, several geographically separate refugia were present within the Iberian refugium (Gomez & Lunt 2007; Ferrero *et al.* 2011). The strong genetic differentiation of a large number of Iberian species into a western (Atlantic) and eastern (Mediterranean) lineages is thought to reflect the disjunct LGM distribution of the most favourable climatic conditions in the peninsula (Schmitt 2007) and the harsher climate of the central Iberian plateau that separates them (Gomez & Lunt 2007).

Unlike other bat species for whom Iberia was the principal glacial refugia (e.g. *Plecotus austriacus*; Razgour *et al.* 2013), *M. escalerai* is unique as it has never expanded its range beyond the peninsula, even though it is found across the Pyrenees, and therefore, Iberia for this species may represent an area of endemism rather than refugium (Stewart *et al.* 2010). Other Iberian endemics, like *Galemys pyrenaicus*, show similar patterns of divergence into distinct evolutionary lineages, suggesting the existence of complex isolation mechanisms as species experienced whole glacial processes of contraction and dispersal within the peninsula (Igea *et al.* 2013).

Approximate Bayesian computation model-based inference confirms the presence of separate western and eastern refugia during the LGM and has identified the source populations of each geographical group as colonies that experienced suitable climatic conditions during the LGM based on SDM projections. Moreover, in line with SDM projections of climatic suitability during the LIG, evolutionary history inference suggests that the western group was the source population. The concordance between the projected distribution of suitable climatic conditions during the LGM and LIG based on SDMs and evolutionary history inference based on genetic data lends support to the presence of niche conservatism in climatic tolerance in *M. escalerai*. Niche conservatism may limit the ability of species to adapt to novel environmental conditions within the time frame required to respond to climate changes, suggesting that instead species will either shift their geographic ranges to track suitable climatic regimes or go extinct (Wiens & Graham 2005). However, Pellissier *et al.* (2013) show that, at least for arctic-alpine plant species, niche conservatism is more pronounced at cold than warm thermal limits because biotic interactions (e.g. competition) play a more important role when conditions are less severe and species are not at their physiological limits.

Yet, climate and topography alone do not determine a species' occurrence, as is evident from the full SDM, in which habitat variables, and in particular the presence of coniferous woodlands, was a strong determinant of occurrence probability. Predicted distribution of

forest tree species in Iberia during the LGM (Benito Garzón *et al.* 2007) and evidence from pollen records (Gomez & Lunt 2007; Lopez de Heredia *et al.* 2007; Rodriguez-Sanchez *et al.* 2010) suggest that most of the studied colonies were located in areas where forests persisted during the LGM. Therefore, forest availability is not likely to have been a major limiting factor for *M. escalerai* during colder periods. Although LGM forests were dominated by pines (Rodriguez-Sanchez *et al.* 2010), the main woodland type where *M. escalerai* is currently found based on the full SDM, south and southwestern Iberia was less forested and dominated by evergreen oaks (Benito Garzón *et al.* 2007). This may explain the ABC inference that *M. escalerai* persisted during the LGM in western and eastern areas, while the south was only colonized around the early- to mid-Holocene when the predicted distribution of pines extended to the southwest (Benito Garzón *et al.* 2007).

#### *Current patterns of genetic variability and future losses*

Population assignment and geographical separation was less clear at the microsatellite than the mtDNA level. Only a slight signature of a geographical west and east divide was evident, most colonies were assigned to more than one genetic population cluster, and many individuals showed some level of admixture. Moreover, colony assignment into geographical groups did not always follow the same pattern as the mtDNA data set. For example, based on the mtDNA data set, the north-western colony Ourense belongs to the north-central-eastern lineage, despite being geographically close to one of the western colonies, while the microsatellite data set groups Ourense with the western colonies. This inconsistency in population assignment may reflect the effect of recent (post-LGM) gene flow disguising older population splits. Microsatellites with their higher evolutionary rates reflect recent or contemporary genetic patterns, while mtDNA is more informative of events that occurred during earlier periods of the species' history (Wan *et al.* 2004).

Alternatively, more limited population structure at the microsatellite level may be the result of male-biased dispersal and female philopatry, a common pattern in bat species (Burland & Worthington Wilmer 2001). Ruedi & Castella (2003) identified a similar pattern in *Myotis myotis*, attributing the absence of population structure at the microsatellite level vs. the strong population structure and limited gene flow between colonies at the mtDNA level to the estimated male bias in the proportion of dispersing individuals (>90%). These disparities highlight the importance of combining biparentally inherited nuclear markers and maternally

inherited mtDNA markers with different evolutionary rates in phylogeographic studies.

Genetic diversity, based on both the mtDNA and microsatellite data sets, is highest in southern colonies, despite their more recent evolutionary history based on the ABC inference. Although this region contains several unique haplotypes and private alleles, high levels of genetic diversity may also be due to this region acting as a 'hybrid/contact zone' between the western and eastern refugia, in which genetic diversity was enriched by the admixture of divergent lineages (Hewitt 2001). And indeed, southern colonies include haplotypes that group with both the western and north-central-eastern clades. On the other hand, high levels of inbreeding and reduced allelic diversity in the most northeastern colony (Girona) and the island colony (Illes Balears) may reflect their geographic isolation and limited recent gene flow from other populations. In the northeastern colony in particular, high coancestry values likely reflect inbreeding in a small isolated population, rather than relatedness due to natal philopatry and the presence of mothers and their pups, because this is the only location where samples were collected from a swarming site and not a maternity colony. While bat summer maternity colonies can include a high proportion of relatives due to strong female natal philopatry, during the autumn, the closely related *M. nattereri* tends to migrate away from summer roosts to swarming sites that serve large catchment areas of up to 60 km (Rivers *et al.* 2006).

Under future climate change projections, the relative occurrence probability of *M. escalerai* across most of Iberia is predicted to decrease substantially. Range losses are predicted to be greatest in the south, placing the entire southern lineage in climatically unsuitable areas by 2070. Although low levels of population differentiation between southern colonies and both western and north-central-eastern colonies indicate the presence of gene flow under current conditions, range fragmentation is likely to increase in the future, resulting in colony isolation. Increased isolation can limit future range shifts and lead to increased inbreeding and loss of genetic diversity (Frankham 1995). Future climate change poses a particular threat to *M. escalerai* because it is restricted to the Iberian Peninsula where changes are predicted to be particularly severe (EEA 2012). Other drivers of environmental change, and in particular anthropogenic habitat loss, may hamper the ability of low dispersal and habitat specialist species, like *M. escalerai*, to shift their ranges in response to climate changes (Warren *et al.* 2001).

Forests are predicted to show a time lag in their response to climate change at their trailing edge. Increased temperatures and frequency of droughts are

predicted to reduce seedling recruitment and forest regeneration, but adult trees may be able persist in climatically unsuitable areas due to their longevity and phenotypic plasticity (Jump *et al.* 2009). Because forests provide cooler microclimates that can help buffer the effects of macroclimatic warming (De Frenne *et al.* 2013), *M. escalerai* colonies may be able to persist in climatically unsuitable areas in the short-term owing to their association with forests. Yet, in the longer term, modelling studies predict severe range contractions of mountain conifer, Mediterranean and sub-Mediterranean forests in central and southern parts of the Iberian Peninsula (Benito Garzón *et al.* 2008).

#### *What restricts an endemic species*

Species distribution models predict that suitable areas for *M. escalerai* are available outside the Iberian Peninsula, in particular along the Mediterranean coast of France. Although only limited genetic sampling has taken place so far, both Salicini *et al.* (2013) and Puechmaille *et al.* (2012) genetically identified all samples beyond the eastern French Pyrenees as its congeners *Myotis spA* or *M. nattereri ss*. Yet, more extensive sampling is needed in areas outside the Iberian Peninsula identified by our SDMs as potentially suitable. Individual samples from around the Pyrenees, including the French Pyrenees, fell within the north-central-eastern clade and mostly belonged to the common eastern haplotype, suggesting that this area was colonized from the eastern refugia, rather than form a putative 'northern refugia' (Stewart & Lister 2001).

The range of *M. escalerai* is at least partly restricted by geographical barriers, the Gibraltar Straits in the south and the Pyrenees mountain range in the north (Salicini *et al.* 2013), although the Pyrenees themselves do not appear to form a barrier (Evin *et al.* 2009; Puechmaille *et al.* 2012). The Iberian Peninsula is home to several other restricted range endemic species, whose limited dispersal abilities prevent them from crossing these geographical barriers, and as a result their entire evolutionary history took place within Iberia (e.g. Igéa *et al.* 2013). Although flight offers bats greater vagility, the Pyrenees have formed a geographical barrier for several bat species, restricting both postglacial range recolonization from this refugium (Rebelo *et al.* 2012; Dool *et al.* 2013) and current patterns of gene flow (Razgour *et al.* 2014). Similarly, the Gibraltar Straits delimit the range of several bat species despite their relative narrow breadth (García-Mudarra *et al.* 2009).

However, because *M. escalerai* is found across the Pyrenees, including the French side of the eastern Pyrenees (Evin *et al.* 2009; Puechmaille *et al.* 2012), ecological barriers as a result of biotic interactions, rather than

geographical barriers, may have played a more important role. Interspecific competition with its cryptic congeners *M. spA* and *M. nattereri s.s.* that may occupy similar ecological niches across the rest of Europe (Salicini *et al.* 2013) could have limited the spread of *M. escalerai* beyond the Pyrenees. It is possible that a delay in northward population expansion post-LGM due to the longer persistence of ice cover in the Pyrenees meant that advancing competing congeners from the Italian and Balkan refugia restricted the space available for *M. escalerai* north of the Pyrenees, as has been postulated for some Iberian forest tree lineages (Rodríguez-Sánchez *et al.* 2010). This suggests that future range gains predicted around the north coast of Iberia, where *M. escalerai* is sympatric with *M. spA*, and in western France, north of the Pyrenees, where *M. nattereri s.s.* is present, may not help offset extensive range losses in the south and centre of Iberia because competitive exclusion may limit northern population expansion. However, the presence of some altitudinal segregation in sympatric localities (Agirre-Mendi & Ibáñez 2012) could indicate different ecological optima for each of these two species, which may allow them to coexist in areas of range overlap.

#### **Conclusions**

A concentration of high genetic diversity and deeply differentiated evolutionary lineages in Iberia has been found in other European bat species with limited long-distance dispersal abilities (Ibáñez *et al.* 2006; Dool *et al.* 2013; Razgour *et al.* 2013), highlighting the evolutionary importance of this peninsula for European bats. Here, we resolve the spatial genetic history of a species for which Iberia is not only a glacial refugium but also its range limit, and therefore, its future survival prospects are closely tied to climatic processes occurring within the peninsula. We show that past climatic oscillations resulted in the divergence of *M. escalerai* into separate western and north-central-eastern populations, supporting the 'refugia within refugia' hypothesis. In accordance with other studies of Iberian reptiles and mammals, our ABC model-based inference and paleo-SDMs indicate that the western population is the older, source population. Although contemporary gene flow may mask historic lineage splits, a signature of geographical population structure is still maintained. The role of ecological barriers due to interspecific competition in restricting *M. escalerai* to the Iberian Peninsula, even when climatic conditions are suitable elsewhere, suggest that this species may struggle to shift its range north of the Pyrenees in the future when most of the peninsula is predicted to become climatically unsuitable.

*Myotis escalerae* was identified morphologically a century ago (Cabrera 1904), but because its species status was only recently genetically confirmed (Ibáñez et al. 2006), its global conservation status is yet to be formally assessed, although within Portugal, it is listed as vulnerable (ICNF 2013). Our findings suggest that conservation management for this species should increase landscape connectivity across Iberia in order to facilitate northwestern range shifts in response to future climate change, especially from the southern lineage that is particularly threatened by future changes.

## Acknowledgements

We are grateful for the invaluable support provided by M. Bertozzi, J.L. García-Mudarra and J. Nogueras in the field and laboratory, as well as to all the people who helped with the collection of samples: H. Rebelo, H. Santos, T. Castelló, O. de Paz, D. Garcia, J. Quetglas, X. Puig, C. Flaquer, A. López-Bauccells, G. Schreur, R. Hermida and F. Lamas. Logistical support was provided by Laboratorio de Ecología Molecular, Estación Biológica de Doñana, CSIC (LEM-EBD) and the ISPRA Conservation Genetics Laboratory staff and in particular A. Viglino. We thank the editor, Rémy Petit, John Stewart and three anonymous reviewers for their invaluable comments that greatly improved the manuscript. I. Salicini benefited from a JAE (Junta para la Ampliación de Estudios' programme) pre-doctoral fellowship from the Consejo Superior de Investigaciones Científicas (CSIC). The study was funded by projects 200430E330 of the CSIC, SAF2006-12784-C02-02 of the Spanish Ministry of Science and Education, PPNN181/2010 of the Spanish Ministry of Environment, and with support of the Spanish Severo Ochoa Program (SEV-2012-0262). O. Razgour was funded by the University of Stirling Impact Fellowship.

## References

- Agirre-Mendi PT, Ibáñez C (2012) Primeros datos sobre la distribución de *Myotis cf. nattereri* y *Myotis escalerae* Cabrera, 1904 (Chiroptera: Vespertilionidae) en la Comunidad Autónoma de La Rioja. *Barbastella*, **5**, 8–11.
- Alvarado-Serrano DF, Knowles LL (2014) Ecological niche models in phylogeographic studies: applications, advances and precautions. *Molecular Ecology Resources*, **14**, 233–248.
- Araújo MB, Nogues-Bravo D, Diniz-Filho JAF et al. (2008) Quaternary climate changes explain diversity among reptiles and amphibians. *Ecography*, **31**, 8–15.
- Avise JC (2000) *Phylogeography: The History and Formation of Species*. Harvard University Press, Massachusetts, USA.
- Benito Garzón M, Sanchez de Dios R, Sainz Ollero H (2007) Predictive modelling of tree species distributions on the Iberian Peninsula during the Last Glacial Maximum and Mid-Holocene. *Ecography*, **30**, 120–134.
- Benito Garzón M, Sanchez de Dios R, Sainz Ollero H (2008) Effects of climate change on the distribution of Iberian tree species. *Applied Vegetation Science*, **11**, 169–178.
- Burland TM, Worthington Wilmer J (2001) Seeing in the dark: molecular approaches to the study of bat populations. *Biology Reviews*, **76**, 389–409.
- Cabrera A (1904) Ensayo monográfico sobre los quirópteros de España. *Memorias de la Real Sociedad Española de Historia Natural*, **2**, 249–287.
- Castella V, Ruedi M (2000) Characterization of highly variable microsatellite loci in the bat *Myotis myotis* (Chiroptera: Vespertilionidae). *Molecular Ecology*, **9**, 1000–1002.
- Cornuet JM, Ravigné V, Estoup A (2010) Inference on population history and model checking using DNA sequence and microsatellite data with the software DIYABC (v1.0). *BMC Bioinformatics*, **11**, 401.
- Cornuet J-M, Pudlo P, Veysier J et al. (2014) DIYABC v2.0: a software to make Approximate Bayesian Computation inferences about population history using Single Nucleotide Polymorphism. DNA sequence and microsatellite data. *Bioinformatics*, **30**, 1187–1189.
- Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods*, **9**, 772.
- De Frenne P, Rodriguez-Sánchez F, Coomes DA et al. (2013) Microclimate moderates plant population responses to macroclimate warming. *Proceedings of the National Academy of Sciences of the USA*, **110**, 18561–18565.
- Dool SE, Puechmaille SJ, Dietz C et al. (2013) Phylogeography and postglacial recolonization of Europe by *Rhinolophus hipposideros*: evidence from multiple genetic markers. *Molecular Ecology*, **22**, 4055–4070.
- Earl DA, von Holdt BM (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources*, **4**, 359–361.
- EEA (2012) *Climate change, impacts and vulnerability in Europe 2012: an indicator-based report*. EEA Report no. 12/2012. EEA, Copenhagen. Available at: <http://www.eea.europa.eu/publications/climate-impacts-and-vulnerability-2012>.
- Evin A, Lecoq V, Durand M-O, Tillon L, Pons J-M (2009) A new species for the French bat list: *Myotis escalerae* (Chiroptera: Vespertilionidae). Mammalia, **73**, 142–144.
- Ferrero ME, Blanco-Aguiar JA, Lougheed SC et al. (2011) Phylogeography and genetic structure of the red-legged partridge (*Alectoris rufa*): more evidence for refugia within the Iberian glacial refugium. *Molecular Ecology*, **20**, 2628–2642.
- Frankham R (1995) Effective population size / adult population size ratios in wildlife: a review. *Genetical Research*, **66**, 95–107.
- García-Mudarra JL, Ibáñez C, Juste J (2009) The straits of Gibraltar: barrier or bridge to Ibero-Moroccan bat diversity? *Biological Journal of the Linnean Society*, **96**, 434–450.
- Gómez A, Lunt DH (2007) Refugia within refugia: patterns of phylogeographic concordance in the Iberian Peninsula. Ch. 5. (eds Weiss S, Ferrand N), pp. 155–188. *Phylogeography of Southern European Refugia: Evolutionary Perspectives on the Origins and Conservation of European Biodiversity*. Springer, Dordrecht, the Netherlands.
- Goudet J (1995) FSTAT (Version 1.2): a computer program to calculate F-statistics. *Journal of Heredity*, **86**, 485–486.
- Hampe A, Petit R (2005) Conserving biodiversity under climate change: the rear edge matters. *Ecology Letters*, **8**, 461–467.
- Hansen MC, Potapov PV, Moore R et al. (2013) High resolution global maps of 21st century forest cover change. *Science*, **342**, 850–853.

- Hewitt GM (2000) The genetic legacy of the Quaternary ice ages. *Nature*, **405**, 907–913.
- Hewitt GM (2001) Speciation, hybrid zones and phyogeography – or seeing genes in space and time. *Molecular Ecology*, **10**, 537–549.
- Hewitt GM (2004) Genetic consequences of climatic oscillations in the Quaternary. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **359**, 183–195.
- Hickerson MJ, Carstens BC, Cavender-Bares J et al. (2010) Phylogeography's past, present, and future: 10 years after Avise, 2000. *Molecular Phylogenetics and Evolution*, **54**, 291–301.
- Hofreiter M, Stewart JR (2009) Ecological change, range fluctuations and population dynamics during the Pleistocene. *Current Biology*, **19**, R584–R594.
- Hollingsworth E, Brahana V, Inlander E, Slay M (2008) Karst regions of the world (KROW): global karst datasets and maps to advance the protection of karst species and habitats worldwide. USGS Scientific Investigation Report 2008–5023. Available at: <http://pubs.usgs.gov/sir/2008/5023/06holings.htm>.
- Ibáñez C, Garcia-Mudarra JL, Ruedi M, Stadelmann B, Juste J (2006) The Iberian contribution to cryptic diversity in European bats. *Acta Chiropterologica*, **8**, 277–297.
- ICNF (2013) *Atlas dos Morcegos de Portugal Continental*. Instituto Conservação da Natureza e das Florestas, Portugal.
- Igea J, Aymerich P, Fernandez-Gonzalez A et al. (2013) Phylogeography and postglacial expansion of the endangered semi-aquatic mammal *Galemys pyrenaicus*. *BMC Evolutionary Biology*, **13**, 115.
- IPCC (2013) Climate Change 2013: The Physical Science Basis. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (eds Stocker TF, Qin D, Plattner G-K et al.). Cambridge University Press, Cambridge, UK and New York, USA.
- Jump AS, Mátyás C, Peñuelas J (2009) The altitude-for-latitude disparity in the range retractions of woody species. *Trends in Ecology and Evolution*, **24**, 694–701.
- Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology*, **16**, 1099–1106.
- Kerth G, Safi K, König B (2002) Mean colony relatedness is a poor predictor of colony structure and female philopatry in the communally breeding Bechstein's bat (*Myotis bechsteinii*). *Behavioural Ecology and Sociobiology*, **52**, 203–210.
- Librado P, Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, **25**, 1451–1452.
- Liu C, White M, Newell G (2013) Selecting thresholds for the prediction of species occurrence with presence-only data. *Journal of Biogeography*, **40**, 778–789.
- Lopez de Heredia U, Carrion JS, Jimenez P, Collada C, Gil L (2007) Molecular and palaeoecological evidence for multiple glacial refugia for evergreen oaks on the Iberian Peninsula. *Journal of Biogeography*, **34**, 1505–1517.
- Merow C, Smith MJ, Silander JA (2013) A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography*, **36**, 1058–1069.
- Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, **6**, 288–295.
- Pellissier L, Bräthen KA, Vitzoz P et al. (2013) Thermal niches are more conserved at cold than warm limits in arctic-alpine plant species. *Global Ecology and Biogeography*, **22**, 933–941.
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modelling of species geographic distributions. *Ecological Modelling*, **190**, 231–259.
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics*, **155**, 945–959.
- Puechmaille SJ, Allegrini B, Boston ESM et al. (2012) Genetic analyses reveal further cryptic lineages within the *Myotis nattereri* species complex. *Mammalian Biology*, **77**, 224–228.
- Raes N, ter Steege H (2007) A null-model for significance testing of presence-only species distribution models. *Ecography*, **30**, 727–736.
- Raymond M, Rousset F (1995) GENEPOP (v1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity*, **86**, 248–249.
- Razgour O, Juste J, Ibáñez C et al. (2013) The shaping of genetic variation in edge-of-range populations under past and future climate change. *Ecology Letters*, **16**, 1258–1266.
- Razgour O, Rebelo H, Puechmaille SJ et al. (2014) Scale-dependent effects of landscape variables on gene flow and population structure in bats. *Diversity and Distributions*, **20**, 1173–1185.
- Rebelo H, Froufe E, Brito JC et al. (2012) Postglacial colonization of Europe by the barbastelle bat: agreement between molecular data and past predictive modelling. *Molecular Ecology*, **21**, 2761–2774.
- Rivers NM, Butlin RK, Altringham JD (2006) Autumn swarming behaviour of Natterer's bats in the UK: population size, catchment area and dispersal. *Biological Conservation*, **127**, 215–226.
- Rodriguez-Ramilo ST, Wang J (2012) The effect of close relatives on unsupervised Bayesian clustering algorithms in population genetic structure analysis. *Molecular Ecology Resources*, **12**, 873–884.
- Rodriguez-Sanchez F, Hampe A, Jordano P, Arroyo J (2010) Past tree range dynamics in the Iberian Peninsula inferred through phylogeography and palaeodistribution modelling: a review. *Review of Palaeobotany and Palynology*, **162**, 507–521.
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, **19**, 1572–1574.
- Rosenberg NA (2004) DISTRUCT: a program for the graphical display of population structure. *Molecular Ecology Notes*, **4**, 137–138.
- Rousset F (2008) GENEPOP'007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources*, **8**, 103–106.
- Ruedi M, Castella V (2003) Genetic consequences of the ice ages on nurseries of the bat *Myotis myotis*: a mitochondrial and nuclear survey. *Molecular Ecology*, **12**, 1527–1540.
- Salicini I, Ibáñez C, Juste J (2011) Multilocus phylogeny and species delimitation within the Natterer's bat species complex in the Western Palearctic. *Molecular Phylogenetics and Evolution*, **61**, 888–898.
- Salicini I, Ibáñez C, Juste J (2013) Deep differentiation between and within Mediterranean glacial refugia in a flying mam-

- mal, the *Myotis nattereri* bat complex. *Journal of Biogeography*, **40**, 1182–1193.
- Schmitt T (2007) Molecular biogeography of Europe: Pleistocene cycles and postglacial trends. *Frontiers in Zoology*, **4**, 11.
- Stewart JR, Lister AM (2001) Cryptic northern refugia and the origins of the modern biota. *Trends in Ecology and Evolution*, **16**, 608–613.
- Stewart JR, Lister AM, Barnes I, Dalen L (2010) Refugia revisited: individualistic responses of species in space and time. *Proceedings of the Royal Society of London B: Biological Sciences*, **277**, 661–671.
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, **4**, 535–538.
- VanDerWal J, Shoo LP, Graham C, Williams SE (2009) Selecting pseudo-absence data for presence-only distribution modeling: how far should you stray from what you know? *Ecological Modelling*, **220**, 589–594.
- Velo-Anton G, Godinho R, Harris DJ et al. (2012) Deep evolutionary lineages in a Western Mediterranean snake (*Vipera latastei/monticola* group) and high genetic structuring in Southern Iberian populations. *Molecular Phylogenetics and Evolution*, **65**, 965–973.
- Wan Q-H, Wu H, Fujihara T, Fang S-G (2004) Which genetic marker for which conservation genetics issue? *Electrophoresis*, **25**, 2165–2176.
- Wang J (2007) Triadic IBD coefficients and applications to estimating pairwise relatedness. *Genetical Research*, **89**, 135–153.
- Wang J (2011) COANCESTRY: A program for simulating, estimating and analysing relatedness and inbreeding coefficients. *Molecular Ecology Resources*, **11**, 141–145.
- Warren MS, Hill JK, Thomas JA et al. (2001) Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature*, **414**, 65–69.
- Warren DL, Glor RE, Turelli M (2010) ENMTools: a toolbox for comparative studies of environmental niche models. *Ecography*, **33**, 607–611.
- Wiens JJ, Graham CH (2005) Niche conservatism: integrating evolution, ecology, and conservation biology. *Annual Review of Ecology, Evolution, and Systematics*, **36**, 519–539.
- Xia X, Xie Z (2001) DAMBE: Data analysis in molecular biology and evolution. *Journal of Heredity*, **92**, 371–373.

---

C.I., I.S. and J.J. designed the study, collected or organized the sample collection, generated the molecular data and contributed to the manuscript. E.R. helped with obtaining the microsatellite data set and information at ISPRA Conservation Genetics Laboratory and contributed to the manuscript. O.R. wrote the manuscript and performed the genetic analysis (mtDNA and microsatellite), species distribution modelling and ABC evolutionary history analysis.

---

## Data accessibility

DNA sequences GenBank Accessions nos: KT365234–KT365283 and KT633874 (Table S1–S2, Supporting information).

Data archived in Dryad (doi: 10.5061/dryad.v1v47): aligned mtDNA sequences for all individuals, nexus input file for MrBayes phylogenetic analysis, MrBayes tree file underlying the phylogeny, Structure input files, Structure results and individual assignments, and MaxEnt location records.

## Supporting information

Additional supporting information may be found in the online version of this article.

**Appendix S1** Approximate Bayesian computation supplementary methods.

**Table S1** Geographic location of *Myotis escalerai* colonies included in the study (geographical coordinates Lat/Long, WGS1984), number of individuals sampled from each colony and their haplotypes.

**Table S2** The geographical location of *Myotis escalerai* individual samples included in the mitochondrial DNA study (geographical coordinates Lat/Long, WGS1984) and their haplotypes.

**Table S3** Genetic diversity at the mitochondrial DNA level for all *Myotis escalerai* sequences divided into the three main geographical groups.

**Table S4** Genetic differentiation at the mitochondrial DNA level between *Myotis escalerai* geographical groups based on  $\theta_{ST}$  values (Chi Square = 678.5,  $P < 0.001$ ).

**Table S5** Genetic differentiation at the microsatellite level between *Myotis escalerai* colonies based on  $F_{ST}$  values.

**Table S6** ABC parameters estimations for the full analysis (Analysis 1) based on Scenario 1.1.

**Table S7** ABC parameters estimations for the Western Group analysis (Analysis 2) based on Scenario 2.1.

**Table S8** ABC parameters estimations for the Eastern Group analysis (Analysis 3) based on Scenario 3.1.

**Table S9** ABC parameters estimations for the demographic history analysis (Analysis 4) based on Scenario 4.3.

**Fig. S1** Results of the STRUCTURE analysis for the *Myotis escalerai* microsatellite dataset.

**Fig. S2** Variables included in the full SDM and their relative contribution to the model in terms of their effect on increasing model gain when used in isolation (size of dark blue relative to red bar) and on decreasing model gain when omitted from the model (extent of reduction in light blue bars relative to the red bar).

**Fig. S3** Variables included in the climate SDM and their relative contribution to the model in terms of their effect on increasing model gain when used in isolation (size of dark blue relative to red bar) and on decreasing model gain when omitted from the model (extent of reduction in light blue bars relative to the red bar).

**Fig. S4** The full species distribution model for *Myotis escalerai* with relative occurrence probability ranging from high in blue to low in yellow.

**Fig. S5** Predictions of the species distribution model for *Myotis escalerai* projected to conditions during the last interglacial per-

iod (LIG ~130 000 ya), with relative occurrence probability ranging from high in blue to low in yellow.

**Fig. S6** Areas where predictions when models are hind-casted to the LGM (Last Glacial Maximum) are affected by variables outside their training range.

**Fig. S7** Areas where predictions when models are fore-casted to 2070 (+8.5rcp scenario) are affected by variables outside their training range.

**Fig. S8** The five scenarios compared in the ABC inference of demographic history.